

# Free Radical Scavenging Activity of *Artemisia vulgaris* Essential Oil

Lok Ranjan Bhatt and Seung Hwa Baek\*

Department of Herbal Resources, Professional Graduate School of Oriental Medicine, Wonkwang University, Iksan 570-749, Korea

**Abstract :** The radical scavenging activity of *Artemisia vulgaris* essential oil was evaluated by 1,1-diphenyl-2-picrylhydrazyl (DPPH) assay in this study. Essential oil exhibited a significant free radical scavenging activity, with the highest activity at 15  $\mu\text{L}/\text{mL}$  concentration. The reaction rate was slow and concentration-dependent.

**Key words :** *Artemisia vulgaris* essential oil, Free radical scavenging activity, Butylated hydroxyanisole,  $\alpha$ -Tocopherol, Concentration-dependent manner.

## Introduction

Plant volatile oils are known to have potential natural agents for food preservation as their effectiveness against a wide range of microorganisms has been well established. Essential oils have been reported to have a range of biological properties. The major properties amongst are their antibacterial, antifungal and antioxidant properties.<sup>1</sup> Recently many essential oils have been qualified as natural antioxidants<sup>2,3,4</sup> and are suggested as a potential substitute of synthetic antioxidants. Furthermore, essential oils and their components are gaining an interest because of their relatively safe status, wide acceptance by consumers, and exploitation for potential multi-purpose functional use.<sup>5,6</sup>

*Artemisia vulgaris* Linn (family: Compositae) is a common shrub found in mountains of Nepal (1,500 - 3,600 m). It is one of the most religious plants in Nepal and is offered in many ritual celebrations. It is also used extensively in spiritual treatment of patient. For example local healers use the foliage in chasing the evils away from the patient's body. Infusion of leaves and flowering tops are used in nervous and spasmodic affections, asthma and diseases of brain.<sup>7</sup> Many *Artemisia* species have a characteristic scent or taste, caused by monoterpenes and sesquiterpenes, which in many cases are the reason for their application in folk medicine.<sup>8</sup> So far a few species of Nepalese plants have been studied for their antioxidant activity.<sup>9,10,11</sup> To our best knowledge, there have been no previous studies regarding the free

radical scavenging activity of *Artemisia vulgaris* essential oil. Therefore, the present study was conducted to identify antioxidant properties of essential oil of this plant.

## Materials and Methods

### 1. Instrument and reagents

Octadecyl-functionalized silica gel, butylated hydroxy anisole,  $\alpha$ -tocopherol and the 1,1-diphenyl-2-picrylhydrazyl (DPPH) were purchased from Sigma. All reagents used, including solvents, were of analytical grade. Recordings were made in a UV-vis Diode Array Spectrophotometer, Hewlett Packard 8453.

### 2. Plant materials

*Artemisia vulgaris* essential oil (steam distillation of the leaves) was obtained from Gharelu Herb Processing Centre, Chapagaun, Lalitpur in Nepal in March 2004. The oil has a pale yellow colour with powerful, fresh-camphoraceous aroma.

### 3. DPPH assay

The ability of the *Artemisia vulgaris* essential oil to donate hydrogen atoms or electrons was measured from the bleaching of purple coloured methanol solution of DPPH.<sup>12</sup> Briefly, 0.5 mL of methanolic solution of 0.24 mM DPPH was mixed with 0.5 mL of *Artemisia* essential oil (in methanol) in different concentrations (5, 10, 15, 20, 25 and 30  $\mu\text{L}/\text{mL}$ ) and reference compounds. The mixture was shaken vigorously and left to stand for 30 min in the dark, and the absorbance was then measured at 517 nm against a blank. Radical scavenging activity was expressed as the percentage of DPPH

\*Correspondence to: Seung Hwa Baek Department of Herbal Resources, Professional Graduate School of Oriental Medicine, Wonkwang University, Iksan 570-749, Korea. Tel: +82-64-850-6225; E-mail: shbaek@wonkwang.ac.kr

elimination after 30 minutes and calculated as follow.

Scavenging ability (%) =  $[A_0 - A_1/A_0] \times 100$ , where  $A_0$  was the absorbance of the control and  $A_1$  was the absorbance in the presence of the test compound. BHA and  $\alpha$ -tocopherol were used as positive controls. Moreover, to understand its kinetic behavior, the decrease in absorbance of essential oil and reference standards was studied at 517 nm for every minute until 90 minutes.

#### 4. Statistical analysis

The data are results of triplicate experiments. Microsoft Excel was used to compute means and standard deviation. Differences among all sample means were determined by analysis of variance (ANOVA) using Origin, version 5 (Micro cal Software, Inc, 1991-97) and were considered significant at  $p < 0.05$ .

### Results and Discussion

Radical scavenging activity and kinetic behavior of different concentrations of essential oil and that of standard antioxidants was studied. Essential oil exhibited a concentration dependent radical scavenging activity, with the highest activity at the concentration of 15  $\mu\text{L}/\text{mL}$  (Fig. 1).


The reaction rate was slow and depended on concentration. However, there was a gradual decrease in scavenging activity with increasing concentration above 15  $\mu\text{L}/\text{mL}$  (Figs. 2 and 3). Antioxidants are believed to intercept the free radical chain of oxidations and to

contribute hydrogen from the phenolic hydroxyl groups themselves; thereby forming stable free radicals which do not initiate or propagate further oxidation of lipids.<sup>13</sup>


Furthermore, the presence of available hydrogen atoms in phenol and/or allylic groups represents a good barrier against the primary oxidative process.<sup>14</sup> The present study revealed that the *A. vulgaris* essential oil is an effective hydrogen donor and a primary antioxidant by reacting with the lipid radical. More than 40 components have been identified from the essential oil of *A. vulgaris*, among which the oxygenated monoterpenes as major components (1,8-cineole, camphor and [alpha]-terpineol) followed by sesquiterpenes and one acyclic non-terpenic compound.<sup>15</sup> Similarly, the leaf oil was found to be rich in 1,8-cineole (2.2-12.2%), [alpha]-thujone (0-11.4%), camphor (15.7-23.1%) and isobomeol (9.3-20.9%).<sup>16</sup> Oxygenated monoterpenes are mainly responsible for the antioxidant property of the plant oils which contain them.<sup>17,18,19</sup> The presence of strongly activated methylene groups in these molecules is probably the reason for this behaviour.<sup>14</sup>

The kinetic classification, according to the time at the steady state, has been reported as rapid <5 minutes, intermediate 5–30 minutes and slow >30 minutes.<sup>20</sup> In this study, the reaction rate of essential oil and BHA was slow while that of  $\alpha$ -tocopherol as intermediate which agreed with previous studies<sup>20,21</sup> showing that the kinetic classification of BHA was slow and  $\alpha$ -tocopherol was intermediate.


The present results revealed that the *A. vulgaris* essential oil possessed a remarkable radical scavenging



**Fig. 1.** Free radical scavenging activity (%) of *Artemisia vulgaris* essential oil and reference compounds at 30 minutes. The results are mean  $\pm$  S.D. of triplicate experiments. Significantly different from the control values:  $p < 0.05$ .



**Fig. 2.** Kinetics of *Artemisia vulgaris* essential oil and reference compounds against the DPPH method with various concentrations



**Fig. 3.** Kinetics of *Artemisia vulgaris* essential oil (15 µL/mL) by DPPH method with compared to reference antioxidants.

activity, and might be effective against diseases caused by over production of free radicals. Since there is no report on the radical scavenging activity of *A. vulgaris* essential oil, our study may be considered as the first report. We hope that the results obtained will provide a starting point for the further investigations especially to evaluate its *in vivo* potential in animal models and different antioxidant mechanisms.

### Acknowledgement

We thank Dr. Lokendra R Sharma and Mr. Puran Pd

Kurmi, Department of Plant Resources, Ministry of Forest & Soil Conservation, Nepal for their help in identification of plant. Thanks are also to Gharelu Herb Processing Centre, Chapagaun, Lalitpur, Nepal for providing the essential oil.

### References

1. Deans, S.G., Waterman, P.G. In: R. K. M. Hay, & P. G. Waterman (Eds.), *Volatile Oil Crops: Their Biology, Biochemistry and Production*. London, Longman, 1993, pp. 113.

2. Aeschbach, R., Löliger, J., Scott, B.C., Murcia, A., Butler, J., Halliwell, B. and Aruoma, O.I. Antioxidant action of thymol, carvacrol, 6-gingerol, zingerone and hydroxytyrosol. *Food Chem. Toxicol.*, 1994, 32, 31-36.
3. Yanishlieva, N.V., Marinova, E.M., Gordon, M.H. and Raneva, V.G. Antioxidant activity and mechanism of action of thymol and carvacrol in two lipid systems. *Food Chem.*, 1999, 64, 59-66.
4. Gianni S., Silvia M., Mariavittoria M., Martina S., Stefano M., Matteo R., Renato, B. Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in foods. *Food Chem.*, 2005, 91, 621-632.
5. Ormancey, X., Sisalli, S., Coutiere P. Formulation of essential oils in functional perfumery, *Parfums, Cosmétiques, Actualités* 2001, 157, 30-40.
6. Sawamura M. Aroma and functional properties of Japanese yuzu (*Citrus junos* Tanaka) essential oil, *Aroma Res.*, 2000, 1, 14-19.
7. Medicinal plants of Nepal, Bulletin of the Department of Medicinal Plants Number 3, His Majesty's Government of Nepal, Ministry of Forest and Soil Conservation, Department of Plant Resources, Thapathali, Kathmandu, Nepal, 1997.
8. Khaled, F. El-Massry, A., El-Ghorab, H., Farouk, A. Antioxidant activity and volatile components of Egyptian *Artemisia judaica* L. *Food Chem.*, 2002, 79, 331-336.
9. KC, S.K., Klaus, M.J. Medicinal plants from Nepal; II. Evaluation as inhibitors of lipid peroxidation in biological membranes. *J. Ethnopharmacol.* 1999, 64, 135-139.
10. Bhandari, M.R., Kawabata, J., Organic acid, phenolic content and antioxidant activity of wild yam (*Dioscorea* spp.) tubers of Nepal. *Food Chem.* 2004, 88(2), 163-168.
11. Krenn, L., Presser, A., Pradhan, R., Bahr, B., Paper, D.H., Mayer, K.K., Kopp, B. Sulfemodin 8-O-beta-D-glucoside, a new sulfated anthraquinone glycoside, and antioxidant phenolic compounds from *Rheum emodi*, *J. Nat. Prod.* 2003, 66, 1107.
12. Yen, G.C., Chen, H.Y. Antioxidant activity of various tea extracts in relation to their antimutagenicity. *J. Agri. Food Chem.*, 1995, 43, 27-37.
13. Dziezak, J.D. Antioxidants. *Food Tech.*, 1986, 40, 94-102.
14. Giuseppe R., Maria T.B. Antioxidant activity of selected essential oil components in two lipid model systems *Food Chem.*, 2000, 69, 167-174.
15. Thau, N.T.P., Thuy, N., Hoi, T.M., Thai, T.H. et al *Artemisia vulgaris* L. from vietnam: chemical variability and composition of the oil along the vegetative life of the plant. *J. Essential Oil Res.*, July/August 2004.
16. Haider, F., Dwivedi, P.D., Naqvi, A.A., Bagchi G.D. Essential oil composition of *Artemisia vulgaris* harvested at different growth periods under Indo-gangetic plain conditions. *J. Essential Oil Res.*, Nov/Dec 2003.
17. Aeschbach, R., Löliger, J., Scott, B.C., Murcia, A., Butler, J., Halliwell, B. and Aruoma, O.U., Antioxidant action of thymol, carvacrol, 6-gingerol, zingerone and hydroxytyrosol. *Food Chem. Toxicol.*, 1994, 32, 31-36.
18. Baratta, M.T., Dorman, H.J.D., Deans, S.G., Biondi, D.M., Ruberto, G. 1998. Chemical composition, antimicrobial and antioxidative activity of laurel, sage, rosemary, oregano and coriander essential oils. *J. Essential Oil Res.*, 1998, 10, 618-627.
19. Laguori, V., Blekas, G., Tsimidou, M., Kokkini, S., Boskou, D. Composition and antioxidant activity of essential oils from oregano plants growth wild in Greece. *Zeitschrift für Lebensmittel-untersuchung und-forzchung* 1993, 197, 20-23.
20. Sanchez-Moreno, C., Larrauri, J., Saura-Calixto, F. A procedure to measure the antiradical efficiency of polyphenols. *J. Sci. Food Agri.*, 1998, 76, 270-276.
21. Xu, J., Chen, S., Hu, Q. Antioxidant activity of brown pigment and extracts from black sesame seed (*Sesamum indicum* L.). *Food Chem.*, 2005, 91, 79-83.